



US006177275B1

(12) **United States Patent**  
**Coruzzi et al.**

(10) **Patent No.:** **US 6,177,275 B1**  
(45) **Date of Patent:** **Jan. 23, 2001**

- (54) **PLANT NITROGEN REGULATORY P-PII GENES**
- (75) Inventors: **Gloria M. Coruzzi**, New York, NY (US); **Hon-Ming Lam**, Hong Kong (HK); **Ming-Hsiun Hsieh**, Woodside, NY (US)
- (73) Assignee: **New York University**, New York, NY (US)
- (\* ) Notice: Under 35 U.S.C. 154(b), the term of this patent shall be extended for 0 days.
- (21) Appl. No.: **08/899,330**
- (22) Filed: **Jul. 23, 1997**

#### Related U.S. Application Data

- (60) Provisional application No. 60/022,328, filed on Jul. 24, 1996.
- (51) **Int. Cl.**<sup>7</sup> ..... **C12N 5/10**; C12N 15/29; C12N 15/82; A01H 5/00
- (52) **U.S. Cl.** ..... **435/468**; 435/69.1; 435/320.1; 435/419; 435/471; 536/23.6; 800/278; 800/298
- (58) **Field of Search** ..... 435/69.1, 410, 435/419, 468, 471, 320.1; 536/23.6; 800/278, 295, 298

#### (56) **References Cited** **PUBLICATIONS**

Stam et al, *Annals of Botany*, vol. 79, pp. 3–12, 1997.\*

Colonna–Romano et al., 1987, “Tight linkage of *glnA* and a putative regulatory gene in *Rhizobium leguminosarum*,” *Nucleic Acids Res.* 15:1951–1964.

De Zamaroczy et al., 1990, “Characterization of three different nitrogen–regulated promoter regions for the expression of *glnB* and *glnA* in *Azospirillum brasilense*,” *Mol. Gen. Genet.* 224:421–430.

Edwards & Coruzzi, 1989, “Photorespiration and light act in concert to regulate the expression the nuclear gene for chloroplast glutamine synthetase,” *Plant Cell* 1:241–248.

Faure et al., 1994, “*Zea3*: A pleiotropic mutation affecting cotyledon development, cytokinin resistance and carbon–nitrogen metabolism,” *Plant J.* 5:481–491.

Foor et al., 1975, “Regulation of synthesis of glutamine synthetase by adenylylated glutamine synthetase,” *Proc. Natl. Acad. Sci. USA* 72:4844–4848.

Forchhammer & De Marsac, 1994, “The P<sub>II</sub> protein in the cyanobacterium *Synechococcus* sp. strain PCC 7942 is modified by serine phosphorylation and signals the cellular N–status,” *J. Bacteriol.* 176:84–91.

Holtel & Merrick, 1988, “Identification of the *Klebsiella pneumoniae glnB* gene: Nucleotide sequence of wild–type and mutant alleles,” *Mol. Gen. Genet* 215:134–138.

Kranz et al., 1990, “Inactivation, sequence, and *lacZ* fusion analysis of a regulatory locus required for repression of nitrogen fixation genes in *Rhodobacter capsulatus*,” *J. Bacteriol.* 172:53–62.

Lam et al., 1994, “Metabolic regulation of the gene encoding glutamine dependent asparagine synthetase in *Arabidopsis thaliana*,” *Plant Physiol.* 106:1347–1357.

B. Magasanik, 1988, “Reversible phosphorylation of an enhancer binding protein, regulates the transcription of bacterial nitrogen utilization genes,” *TIBS* 13:475–479.

B. Magasanik, 1993, “The regulation of nitrogen utilization in enteric bacteria,” *J. Cell. Biochem.* 51:34–40.

Martin et al., 1989, “*Bradyrhizobium japonicum glnB*, a putative nitrogen–regulatory gene, is regulated by NtrC at tandem promoters,” *J. Bacteriol.* 171:5638–5645.

Ninfa & Magasanik, 1986, “Covalent modification of the *glnG* product, NR<sub>1</sub>, by the *glnL* product, NR<sub>2</sub>, regulates the transcription of the *glnALG* operon in *Escherichia coli*,” *Proc. Natl. Acad. Sci. USA* 85:5909–5913.

Son & Rhee, 1987, “Cascade control of *Escherichia coli* glutamine synthetase,” *J. Biol. Chem.* 262:8690–8695.

Souillard & Sibold, 1989, “Primary structure, functional organization and expression of nitrogenase structural genes of the thermophilic archaeobacterium *Methanococcus thermolithotrophicus*,” *Mol. Microbiol.* 3:541–551.

C. Stewart, 1993, “The powers and pitfalls of parsimony,” *Nature* 361:603–607.

Tsinoremas et al., 1991, “Photosynthetic electron transport controls nitrogen assimilation in cyanobacteria by means of posttranslational modification of the *glnB* gene product,” *Proc. Natl. Acad. Sci. USA* 88:4565–4569.

van de Loo et al., 1995, “Expressed sequence tags from developing castor seeds,” *Plant Physiol.* 108:1141–1150.

van Heeswijk et al., 1993, “The genes of the glutamine synthetase adenylation cascade are not regulated by nitrogen in *Escherichia coli*,” *Mol. Microbiol.* 9:443–457.

Vincentz et al., 1993, “Regulation of nitrate and nitrite reductase expression in *Nicotiana plumbaginifolia* leaves by nitrogen and carbon metabolites,” *Plant J.* 3:315–324.

Wray et al., 1994, “The nitrogen–regulated *Bacillus subtilis nrgAB* operon encodes a membrane protein and a protein highly similar to the *Escherichia coli glnB*–encoded P<sub>II</sub> protein,” *J. Bacteriol.* 176:108–114.

\* cited by examiner

*Primary Examiner*—Gary Benzion  
*Assistant Examiner*—Ashwin D. Mehta

(74) *Attorney, Agent, or Firm*—Pennic & Edmonds LLP

#### (57) **ABSTRACT**

The present invention generally relates to plant nitrogen regulatory PII gene (hereinafter P–PII gene), a gene involved in regulating plant nitrogen metabolism. The invention provides P–PII nucleotide sequences, expression constructs comprising said nucleotide sequences, and host cells and plants having said constructs and, optionally expressing the P–PII gene from said constructs. The invention also provides substantially pure P–PII proteins.

The P–PII nucleotide sequences and constructs of the inventor may be used to engineer organisms to overexpress wild–type or mutant P–PII regulatory protein. Engineered plants that overexpress or underexpress P–PII regulatory protein may have increased nitrogen assimilation capacity. Engineered organisms may be used to produce P–PII proteins which, in turn, can be used for a variety of purposes including in vitro screening of herbicides. P–PII nucleotide sequences have additional uses as probes for isolating additional genomic clones having the promoters of P–PII gene. P–PII promoters are light– and/or sucrose–inducible and may be advantageously used in genetic engineering of plants.




FIG. 1A




FIG. 1B




FIG.2






FIG.3



1 2  
FIG.4A



1 2 3 4 5  
FIG.4B




FIG.5




FIG.6




FIG. 7






FIG.8




FIG.9




FIG.10




FIG.11

1 ctgaaagttg tgtaaataaa aaaactagaa tcatggcggc gtcaatgacg  
51 aaacccatct caataacttc tctcggtttc tattctgacg gaaagaacat  
101 tgctttctct gattgcattt cgatttggtc tggattcaga cattcccgac  
151 catcttgctc cgatttggtc acaaagtcac cgagtaataa cagtcgtggt  
201 ttacctgtcg ttagtgccca aatatcttct gattatattc cagactcgaa  
251 attttacaag gtggaagcaa ttgtcagacc atggagaatc cagcaagttt  
301 catcggcttt actgaaaatc gggattcgag gtgttactgt ttctgatgtg  
351 agagggtttg gtgcacaagg aggttctacc gagagacacg gtggctctga  
401 gttctcggaa gacaaatttg ttgctaaagt taagatggaa atcgttgtta  
451 agaaagacca agtggaatct gtaatcaaca caataattga aggagcaagg  
501 acaggagaga ttggtgatgg caagattttt gttttgcctg tgtcagatgt  
551 cataagagtt aggacaggtg agcgtgggga gaaagcagag aagatgactg  
601 gtgatatgct ttcaccgtct taggaacaaa cagagctcaa gaatggtttt  
651 tttttttttc atttcggctc ctagattctg cgaataataa tgaatggagt  
701 ctgtgtttgg tttcatgttg aatcgatcaa gatgtgtttt taactgtaca  
751 tgaattatgc agaaacatct gtcttggttc tcagacatcg aaactctggt  
801 cctaataaaa aaaaaaa

FIG. 12

1 GCGGTGTCGG CCGCTCTAGA ACTAGTGGAT CCCCCGGGCT GCAGGAATTC  
51 GGCACGAGGC TACTGCGAAA CTGGGCTTGC TCACTCCTCT TCATTCTAAT  
101 AACATCAAGA AAGAATTCCC TGTTTTTGAT TTCAGTTTGT TTTGTCCAGA  
151 GCTTAGACAT TCTCGGTTTT CTCACITTTAA CACCGCGGTC AAGCGCGTAA  
201 GATATGCCCC CGTCGTTTCT GTGATTAATG CCCAAAGCTC GCCTGACTAC  
251 ATTCCTGATG CTAAATTCTA CAAAGTGGAA GCAATTCTCA GGCCCTGGCG  
301 AGTCTCGCAA GTTTCCTCGG CTTTGCTAAA AATTGGTATT CGAGGTGTTA  
351 CTGTTTCTGA TGTTTCGAGGT TTTGGTGCTC AAGGTGGTTC AACTGAGAGG  
401 CAGGGCGGCT CAGAATTTTC TGAAGACAAG TTTGTTGCTA AAGTTAAGAT  
451 GGAGATCGTG GTTAGCAAAG ACCAGGTTGA GGATGTTATA GAAAAAATCA  
501 TTGAGGAGGC AAGAACTGGA GAGATTGGAG ACGGCAAGAT TTTCTTGCTG  
551 CCTGTTTCAG ATGTAATAAG AGTCCGCACT GGTGAGCGGG GTGATAAGGC  
601 TGAGAGGATG ACAGGAGGGC GATCTGACAT GAGTACTTCT GCTTGACTGC  
651 TGTGACCAGC AATATAGCAT TCAGGACTAA CTGTCCTTTG AGAAAGCCCC  
701 GCCCTTATTA GCCATTATCC AGTATAGCTT GATAATTTGA ATTTTTTGT  
751 TTCTTAACTA AAGAAACAAA GATCTTTTCA TTATCCTGTT GATGATAATT  
801 GAAAACGGAA GGATCGCGAA TTTGTTCAAG TGCTTGCAAG ATAAATAACA  
851 AGAAGAGGAG TAATGTTAAC AAAAAAAAAA AAAAAAAAAA ACTCGAG

FIG. 13

## PLANT NITROGEN REGULATORY P-PII GENES

This application claims benefit to provisional Application No. 60/022,328 filed Jul. 24, 1996.

This invention was made with U.S. government support under National Institute of Health grant number GM32877 and Department of Energy grant number DE-FG02-94ER20133 and National Science Foundation grant No. DIR-8908095. The U.S. government has certain rights in the invention.

### 1. INTRODUCTION

The present invention generally relates to a plant nitrogen regulatory PII gene (hereinafter P-II gene), a gene involved in regulating plant nitrogen metabolism. The invention provides P-II nucleotide sequences, expression constructs comprising said nucleotide sequences, and host cells and plants having said constructs and, optionally expressing the P-II gene from said constructs. The invention also provides substantially pure P-II proteins.

The P-II nucleotide sequences and constructs of the invention may be used to engineer organisms to overexpress wild-type or mutant P-II regulatory protein. Engineered plants that overexpress or underexpress P-II regulatory protein may have increased nitrogen assimilation capacity. Engineered organisms may be used to produce P-II proteins which, in turn, can be used for a variety of purposes including in vitro screening of herbicides. P-II nucleotide sequences have additional uses as probes for isolating additional genomic clones having the promoters of P-II gene. P-II promoters are light- and/or sucrose-inducible and may be advantageously used in genetic engineering of plants.

### 2. BACKGROUND OF THE INVENTION

Plants can assimilate soil ammonia or nitrate reduced to ammonia into organic form in leaves or roots. Ammonia assimilation into glutamine and glutamate occurs primarily in leaf chloroplasts or in root plastids by the combined action of chloroplast glutamine synthetase (GS2; GLN2 gene) and glutamate synthase (GOGAT) (Miflin, B. J. & Lea, P. J., 1977, *Ann. Rev. Plant Physiol.* 28:299-329). As the assimilation of inorganic nitrogen into organic form requires carbon skeletons, reducing equivalents, and ATP, light serves to coordinate nitrogen assimilation with photosynthesis. Genes involved in plant nitrogen assimilation are induced directly by light (via phytochrome), as well as indirectly by metabolic changes in photosynthate. For example, it has been shown that sucrose supplementation to plant growth media can at least partially induce the expression of mRNA for GLN2 or nitrate reductase (NR) in the absence of light (Cheng et al., 1992, *Proc. Natl. Acad. Sci. USA.* 89:1861-1864; Faure et al., 1994, *Plant J.* 5:481-491). Conversely, sucrose can repress the expression of asparagine synthetase (ASN1) (Lam et al., 1994, *Plant Physiol.* 106:1347-1357). More recently, it has been shown that the effects of sucrose on gene expression can be reversed by the addition of an organic nitrogen source both for nitrate reductase (NR) (Vincentz et al., 1993, *Plant J.* 3:315-324) and for ASN1 (Lam et al., 1994, *Plant Physiol.* 106:1347-1357). These findings indicate that plants are able to sense levels of carbon and organic nitrogen, and in turn modulate the expression of genes involved in nitrogen assimilation.

Bacteria can also assimilate ammonia into glutamate or glutamine. Plants' ability to sense changes in the levels of

carbon and nitrogen metabolites is reminiscent of a nitrogen regulatory system (Ntr) in bacteria in which a protein called PII, encoded by the *glnB* gene, can regulate the assimilation of nitrogen into glutamine via glutamine synthetase (GS; *glnA*) in response to changes in the ratio of organic nitrogen to carbon metabolites (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40).

In response to changes in the metabolic status (i.e., ratio of glutamine to  $\alpha$ -ketoglutarate [*gln*/ $\alpha$ -KG]), the PII protein of bacteria interacts with a set of partners to regulate the *glnA* gene at the transcriptional level, and to regulate GS enzyme activity at the post-translational level (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40). Changes in the *gln*/ $\alpha$ -KG ratio affect the activity of the PII protein via a post-translational modification (uridylylation) at Tyr51 (Magasanik, B., 1988, *TIBS* 13:475-479). In response to low *gln*/ $\alpha$ -KG, the PII protein is uridylylated by uridylyltransferase (UTase) (id.). The PII-UMP thus formed then interacts with an adenylyltransferase (ATase) to deadenylylate the GS-AMP enzyme and thereby activate the GS enzyme (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40). A high *gln*/ $\alpha$ -KG ratio causes the deuridylylation of PII-UMP. This unmodified form of PII interacts with ATase to stimulate the adenylylation and inactivation of the GS enzyme. The ability of ATase to attach or remove AMP from the GS enzyme is dependent on the interaction of ATase with PII or PII-UMP, respectively (Foor et al., 1975, *Proc. Natl. Acad. Sci. USA* 72:4844-4848). Thus, the nitrogen-regulatory protein PII, is a signal transducer whose post-translational modification indirectly regulates GS enzyme activity post-translationally. In addition to its ability to regulate the GS holoenzyme activity, PII can also interact with a two-component system (NRII/NRI, or NtrB/NtrC) to regulate the transcription of the *glnA* gene encoding GS (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40). Under low *gln*/ $\alpha$ -KG levels, NRII-kinase phosphorylates NRI which then interacts with the  $\sigma^{54}$  to activate *glnA* gene expression (Ninfa, A. J. and Magasanik, B., 1986, *Proc. Natl. Acad. Sci. USA* 83:5909-5913). When the *gln*/ $\alpha$ -KG ratio is high, the interaction of PII with NRII stimulates the NRII-phosphatase activity to dephosphorylate NRI-phosphate, and turn off the inducible promoter of *glnA* transcription (Ninfa, A. J. and Magasanik, B., 1986, *Proc. Natl. Acad. Sci. USA* 83:5909-5913). Thus, the nitrogen-regulatory protein PII works in concert with other proteins, including UTase, ATase, NRII, and NRI, to regulate glutamine synthetase enzyme activity or *glnA* transcription in response to the ratio of organic nitrogen to carbon metabolites (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40).

To date, PII homologues have been identified in a diverse set of bacteria including enteric bacteria (Magasanik, B., 1994, *J. Cell. Biochem.* 51:34-40), cyanobacteria (Tsinoremas et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4565-4569), *Bacillus* (Wray et al., 1994, *J. Bacteriol.* 176:108-114), and in archaeobacteria (Souillard, N. and Sibold, I., 1989, *Mol. Microbiol.* 3:541-551).

### 3. SUMMARY OF THE INVENTION

The present invention relates to a plant nitrogen regulatory P-II gene involved in regulating nitrogen assimilation in plants. The invention provides P-II coding nucleotide sequences, expression constructs comprising P-II coding sequences, and host organisms, including plants, containing said expression constructs. The invention also provide P-II proteins.

The invention is based on the surprising discovery that plants have a structural homolog, P-II, to the bacterial PII

protein. This is the first time a PII-like gene has been identified in an eukaryote. The regulation of P-II mRNA levels by light and by metabolites, such as sucrose, parallels those of nitrogen assimilatory genes such as chloroplastic GS2 (GLN2). See Faure et al., 1994, *Plant J.* 5:481-491; Edwards, J. W. and Coruzzi, G. M., 1989, *Plant Cell* 1:241-248; Lam et al., 1994, *Plant Physiol.* 106:1347-1357; Vincentz et al., 1993, *Plant J.* 3:315-324. These findings indicate that like bacterial PII, P-II protein is a plant nitrogen regulatory protein that controls the expression of nitrogen assimilation functions.

The P-II nucleotide sequences and constructs of the invention may be advantageously used to engineer plants to overexpress P-II regulatory protein. P-II overexpression or underexpression should enhance the levels of certain nitrogen assimilation functions and thereby increase nitrogen utilization efficiencies of engineered plants.

P-II nucleotide sequences and constructs of the invention also may be used to engineer organisms to overexpress wild-type or mutant P-II regulatory protein. Full length cDNAs for P-II can be used in a "reverse biochemical" approach to synthesize and characterize the encoded P-II proteins. The ability to use the cloned P-II to synthesize the purified P-II proteins will allow a characterization of P-II protein in terms of physical properties (i.e., inducer or activator preference) and subcellular localization (i.e., plastid vs. cytosol).

Full length P-II cDNA clones may also be used to synthesize highly purified preparations of the wild-type or altered P-II in vitro or in vivo (e.g., bacteria, algae, neurospora, yeast, plant, or animal cells). In vitro or in vivo synthesized P-II protein can be used as a substrate in a screen to identify novel herbicidal compounds, which selectively inhibit this nitrogen regulatory protein. The isolated cDNAs encoding P-II may also be used to create plants resistant to such herbicides.

Nucleic acids, DNA or RNA, encoding P-II have additional uses as probes for isolating additional genomic clones having the promoters of P-II genes. P-II promoters are light- /or sucrose-inducible and may be advantageously used in genetic engineering of plants. For example, P-II promoters may be used to directly express genes encoding functions that P-II proteins directly or indirectly activate or induce. P-II promoter sequences also may be used to construct chimeric promoters.

### 3.1. ABBREVIATIONS AND DEFINITIONS

The following terms as used herein, whether in the singular or plural, shall have the meanings indicated.

Asr=asparagine

AS=asparagine synthetase

bp=basepair

chimeric gene=A chimeric gene comprises a coding sequence linked to a promoter that said coding sequence is not naturally linked to. The coding sequence may encode messenger RNA (mRNA), antisense RNA or ribozymes. As used herein the term is meant to encompass only artificially produced genes using genetic selection or recombinant DNA methodologies and not any genes found in wild-type organisms or cells not subjected to experimentation

EDTA=disodium ethylene diamine tetracetate

GLB1=the gene encoding P-II protein

GS=glutamine synthetase

GUS=1,3- $\beta$ -Glucuronidase

heterologous=unrelated, e.g., not of the same gene or organism

kb=kilobase

operably linked=The linkage between a promoter and a coding sequence such that the promoter controls the transcription of the coding sequence

PCP=polymerase chain reaction

P-II=plant PII

poly(A)=polyadenylate

recombinant=resulting from artificial, genetic-engineering efforts and not natural mutational events

SDS=sodium dodecyl sulfate

20xSSC=175.3 g NaCl and 88.2 g sodium citrate in 800 ml H<sub>2</sub>O, pH 7.0 (adjusted with 10 N NaOH), adjusted to 1 liter.

Peptide and polypeptide sequences defined herein are represented by one-letter symbols for amino acid residues as follows:

A (alanine)

R (arginine)

N (asparagine)

D (aspartic acid)

C (cysteine)

Q (glutamine)

E (glutamic acid)

G (glycine)

H (histidine)

I (isoleucine)

L (leucine)

K (lysine)

M (methionine)

F (phenylalanine)

P (proline)

S (serine)

T (threonine)

W (tryptophan)

Y (tyrosine)

V (valine)

### 4. BRIEF DESCRIPTION OF THE FIGURES

FIG. 1(A-B). Comparison of the deduced amino acid sequences of plant PII and microbial PII polypeptides. FIG. 1A. The amino acid residues conserved between all microbial PIIs and plant PII are shaded. The asterisks (\*) indicate residues conserved between plant and any bacterial PII. Boxed areas labeled I and II include residues of two PII signature domains conserved between all PIIs. The position of the tyrosine residue (Tyr 51) which is uridylylated in *E. coli* is indicated by an arrow. The PII sequences shown in this figure are: *At*, *Arabidopsis thaliana* (SEQ ID NO:1); *Ric*, *Ricinus communis* (SEQ ID NO:2); *Kp*, *Klebsiella pneumoniae* (SEQ ID NO:3); *Ec*, *Escherichia coli* (SEQ ID NO:4); *R1*, *Rhizobium leguminosarum* (SEQ ID NO:5); *Bj*, *Bradyrhizobium japonicum* (SEQ ID NO:6); *Az*, *Azospirillum brasilense* (SEQ ID NO:7); *Rc*, *Rhodobacter capsulatus* (SEQ ID NO:8); *Sy*, *Synechococcus* (SEQ ID NO:9) strain PCC 7942; *Mt1*, *Methanococcus thermolithotrophicus*, glnB-like protein 1 (SEQ ID NO:10) *Mt2*, *Methanococcus thermolithotrophicus* glnB-like protein 2 (SEQ ID NO:11). For best alignment of *Mt2* with glnB, the last two residues, BN, and the peptide sequence FSANLPEIVDIQKII (SEQ ID NO:12) are deleted. This deletion is indicated by "" in the *Mt2* sequence. The numbers 1, 51, and 112 indicate the residue positions of the *E. coli* PII protein. FIG. 1B. The



relationship between plant PII and microbial PIIs is shown by a phylogenetic analysis using parsimony. Bacillus PII-like protein is an anomaly which does not cluster with bacterial taxa. F, phenylalanine; Y, tyrosine.

FIG. 2. Southern blot analysis of GLB1 gene in Arabidopsis genomic DNA. *Arabidopsis thaliana* (Columbia ecotype) genomic DNA was digested with the following restriction enzymes; B, BglII (lane 1); D, DraI (lane 2); N, NdeI (lane 3); Sa, 5Sacl (lane 4); Sm, SmaI (lane 5), resolved by gel electrophoresis, transferred to Hybond-N membrane and probed with DIG-labeled single-stranded probe derived from the Arabidopsis GLB1 cDNA.

FIG. 3. Northern blot analysis of Arabidopsis GLB1 mRNA levels in different organs. Total RNA (20 µg) from roots (lane 1), leaves (lane 2) and flowers (lane 3) was detected with DIG-labeled single-stranded DNA probe derived from Arabidopsis GIB1 cDNA. Roots and leaves were from 6-week-old mature plants. Flowers were buds or developing flowers.

FIG. 4(A-B). Light-induced expression of Arabidopsis GLB1 gene. FIG. 4A. Total RNA (20 µg) from leaves of light-grown (L; lane 1) or dark-adapted (D; lane 2) mature Arabidopsis plants. GLN2 and ASN1 mRNAs were also detected on duplicate Northern blots as controls for RNA preparation. FIG. 4B. Time-course of light-induction of GLB1 mRNA in Arabidopsis. Total RNA (10 µ) from two-week-old Arabidopsis plants grown in semi-hydroponics which were dark-adapted for two days and then exposed to light for 2 hr (lane 1), 4 hr (lane 2), 8 hr (lane 3), 16 hr (lane 4), and 24 hr (lane 5). Arabidopsis DNA probes for GLB1 and GLN2 cDNAs were used to detect mRNAs on the identical Northern blot simultaneously.

FIG. 5. Sucrose-induced expression of Arabidopsis GLB1 gene. Total RNA (10 µg) from two-week-old Arabidopsis plants dark-spotted (lanes 1-3) or light-grown (lanes 4-6) for two days was used for Northern blot analysis. During the dark- or light-treatments, plants were grown in media containing 0% sucrose (lanes 1 and 4), 3% sucrose (lanes 2 and 5) or 3% mannitol (lanes 3 and 6). As a control, 18 S ribosomal RNA was detected on a replicate blot. The ethidium bromide staining of the RNA in the agarose gel before transfer is also shown.

FIG. 6. PII (GLB1) mRNA is induced by light and/or high sucrose. Induction of PII (GLB1) mRNA by light/sucrose mirrors that of the putative downstream target nitrogen assimilatory genes GS2 (GLN2) and Fd-GOGAT (GLU1). PII mRNA is low in dark-adapted plants (lane 1) and is induced by sucrose in the absence of light (lane 2), but not by mannitol, a non-metabolizable sugar (lane 3). Light induces PII mRNA (lane 4). Light plus sucrose super-induces PII mRNA (lane 5). Mannitol cannot superinduce PII mRNA above light levels (lane 6). Nitrogen assimilatory genes GS2 and Fd-GOGAT GLU1, also show induction by sucrose (lane 2) or light (lane 3), but are not super-induced by light and sucrose (lane 5).

FIG. 7. Metabolic control model whereby light or high sucrose induces PII (GLB1) and putative downstream nitrogen assimilatory genes. PII induction by light and/or high sucrose, mirrors the induction of downstream nitrogen assimilatory genes for glutamine synthetase (GLN2), Fd-dependent glutamate synthase (Fd-GOGAT, GLU2), and aspartate aminotransferase (ASP2). By contrast, genes for asparagine synthetase (ASN1) and glutamate dehydrogenase are repressed by high sucrose (Lam et al, 1994 plant physiol. 106: 1347-1357).

FIG. 8. PII (GLB1) expression is repressed by high organic nitrogen, relative to carbon. Sucrose induction of PII

(GLB1) mRNA (lane 1) is repressed by addition of organic nitrogen (lanes 2-4). This regulation of PII mirrors the regulation of the putative downstream nitrogen assimilatory gene for glutamine synthetase (GLN2).

FIG. 9. Metabolic control model whereby dark or high organic nitrogen/low carbon represses PII (GLB1). PII and its putative downstream nitrogen assimilatory genes (GLN2, GLU1 and ASP2) are induced by sucrose (lane 1). Sucrose induction is repressed by addition of organic nitrogen as asparagine (lane 2), glutamine (lane 3) and glutamate (lane 4). By contrast, ASN1 (Lam et al, 1994 plant physiol. 106: 1347-1357). and GDH1 are induced by these conditions.

FIG. 10. Arabidopsis PII (GLB1) gene maps to the top arm of chromosome 4. RFLP and recombinant inbred lines of Arabidopsis to map the PII (GLB1) gene to the top arm of chromosome 4 at position 10.8.

FIG. 11. Transgenic tobacco plants expressing antisense PII mRNA. Antisense Arabidopsis PII was introduced into tobacco by Agrobacterium-mediated DNA transfer. Shown are four independent transgenic tobacco lines which express high levels of PII antisense RNA (lines 1, 2, 5 and 10).

FIG. 12. Nucleotide sequence of Arabidopsis P-II cDNA clone (SEQ ID NO:13).

FIG. 13. Nucleotide sequence of Ricinus Castor Bean P-II cDNA clone (SEQ ID NO:14).

## 5. DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the plant PII (P-II; GLB1) gene. The invention provides nucleotide sequence encoding P-II protein, expression constructs that can be used to express or overexpress P-II, and organisms engineered with P-II expression constructs. The invention also provides; P-II proteins.

In accordance with one aspect of the invention, biologically active P-II may be produced by the cloning and expression of the nucleotide coding sequence for P-II or its functional equivalent in appropriate host cells. Successful expression aid production of purified P-II using the full length cDNA clones described and exemplified herein is significant since this regulatory protein has yet to be isolated or purified from plants.

The synthesis of purified P-II protein via genetic engineering techniques which provide for the expression of full length cDNAs may be utilized to obtain antibodies specific for P-II which will enable the characterization of related proteins and subcellular localizations of P-II in plant cells. The recombinantly produced P-II protein can be advantageously used to screen in vitro for compounds that activates or inhibits the protein's activity. Due to P-II's role in regulating a critical metabolic process, such activators and inhibitors are candidate positive or negative plant growth regulatory compounds, (negative plant growth regulators include herbicides).

The cloned P-II cDNAs expressed by microorganisms can be used to screen and identify new plant growth regulatory compounds and to screen and/or select P-II variants. For example, P-II cDNAs genetically altered and expressed in microorganism can be used to screen and/or select altered P-II proteins which are more sensitive to inducers or constitutively active (i.e., active without the need for light- or sucrose induction). Gene constructs encoding such altered P-II may be used in genetic-engineering to enhance the nitrogen assimilation capabilities of plants.

The invention may be divided into the following stages solely for the purpose of description and illustration: (A)

isolation or generation of the coding sequence for P-PII gene(s); (b) construction of an expression vector which will direct the expression of the P-PII coding sequences; (c) transfection of appropriate hosts which are capable of replicating and expressing the P-PII coding sequences to produce biologically active gene products; and (d) identification and/or purification of the P-PII so produced. Once a transformant or transfectant is identified that expresses high levels of biologically active P-PII, the practice of the invention involves the expansion of that clone and the use of that clone in the production of P-PII, the selection of novel plant growth regulators that effect their action through P-PII, and/or the engineering of transgenic plants.

Another aspect of the invention involves the use of the P-PII promoter to direct the expression of heterologous coding sequences. Light or sucrose activates the P-PII promoter in various organs, including shoots, roots and flower. Thus, this promoter can be used to direct the regulated expression of heterologous gene sequences in appropriate hosts.

The invention is demonstrated herein, by way of examples in which cDNAs of P-PII were prepared, cloned and characterized. P-PII cDNA and genomic clones may be isolated using homologous P-PII sequences obtained from related plants. Various aspects of the invention are described in more detail in the subsections below and in the examples that follow.

#### 5.1. P-PII CODING SEQUENCES

The nucleotide coding sequences for Arabidopsis and castor bean P-PII are depicted in FIGS. 12 and 13, respectively. These nucleotide sequences, or fragments or functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of the P-PII gene product, or functionally active peptides or functional equivalents thereof, in appropriate host cells.

The nucleotide sequences encoding the castor bean P-PII gene and the Arabidopsis P-PII gene can be used to screen cDNA libraries obtained from other plant species to identify further plant P-PII genes. A plant cDNA library may be screened under conditions of reduced stringency, using a radioactively or nonradioactively labeled fragment of the castor bean or the Arabidopsis P-PII clone. Alternatively, the Arabidopsis P-PII or the castor bean P-PII sequences can be used to design degenerate or fully degenerate oligonucleotide probes which can be used as PCR probes or to screen plant cDNA libraries. Alternatively, the probes may be used to screen genomic libraries.

The P-PII nucleotide sequences of the invention include (a) the DNA sequence in FIG. 12 (SEQ ID NO:13); (b) the DNA sequence spanning from nucleotide 33 to 620 as shown in FIG. 12 (SEQ ID NO:15); (c) the DNA sequence in FIG. 13 (SEQ ID NO:14); (d) the DNA sequence spanning from nucleotide 50 to 643 as shown in FIG. 13 (SEQ ID NO:16); (e) any nucleotide sequence that hybridizes to the complement of the DNA sequence shown in FIG. 12 and encodes a functionally equivalent product; and (f) any nucleotide sequence that hybridizes to the complement of the DNA sequence shown in FIG. 13 and encodes a functionally equivalent product. Functional equivalents of the P-PII include naturally occurring P-PII in other plant species, the mutant P-PII whether naturally occurring or engineered. The invention also includes degenerate variants of sequences (a) through (f).

The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore complements of, the nucleotide sequences (a) through (f), in the preceding paragraph. Such hybridization conditions may

be highly stringent or less highly stringent. In instances wherein the nucleic acid molecules are deoxyoligonucleotides ("oligos"), highly stringent conditions may refer, e.g., to washing in 6xSSC/0.5% sodium pyrophosphate at 37° C. (for 17-base oligos), 55° C. (for 20-base oligos), and 60° C. (for 23-base oligos). These nucleic acid molecules may encode or act as P-PII antisense, ribozyme and/or triple helix sequences, useful, for example, in P-PII gene regulation.

The P-PII nucleotide sequences of the present invention also include any nucleotide sequence encoding a plant protein containing the amino acid sequences designated At or RIC in FIG. 1A; and (b) any nucleotide sequence that hybridizes to the complement of the DNA sequences that encode the amino acid sequence designated At or RIC in FIG. 1. The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore complements of the nucleotide sequences (a) and (b). Such hybridization conditions may be highly stringent or less highly stringent. In addition to the P-PII nucleotide sequences described above, full length P-PII cDNA or gene sequences present in the same species and/or homologs of the P-PII gene present in other plant species can be identified and readily isolated, without undue experimentation, by molecular biological techniques well known in the art. The identification of homologs of P-PII in related species can be useful for developing plant model systems for purposes of discovering activators or inhibitors of P-PII to modify P-PII in plants to alter plant nitrogen metabolism. Alternatively, such cDNA libraries, or genomic DNA libraries derived from the organism of interest can be screened by hybridization using nucleotides described herein as hybridization or amplification probes.

Screening can be by filter hybridization, using duplicate filters. The labeled probe can contain at least 15–30 base pairs of the P-PII nucleotide sequence as shown in FIGS. 12 and 13. The hybridization washing conditions used should be of a lower stringency when the cDNA library is derived from an organism different from the type of organism from which the labeled sequence was derived.

Low stringency conditions are well known to those of skill in the art, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions see, for example, Sambrook et al., 1989, *Molecular Cloning, A Laboratory Manual*, Cold Springs Harbor Press, N.Y.; and Ausubel et al., 1989, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y.

Alternatively, the labeled P-PII nucleotide probe may be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. The identification and characterization of plant genomic clones is helpful for designing diagnostic tests and clinical protocols for regulating plant growth rate and nitrogen metabolism. For example, sequences derived from regions adjacent to the intron/exon boundaries of the plant gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g. splice acceptor and/or donor sites), etc., that can be used in diagnostics.

Further, a P-PII gene homolog may be isolated from nucleic acid of the organism of interest by performing PCR using two degeneration oligonucleotide primer pools designed on the basis of amino acid sequences within the P-PII gene product disclosed herein. The template for the reaction may be cDNA obtained by reverse transcription of mRNA prepared from, for example, plant cell lines or tissue, known or suspected to express P-PII gene allele.

The PCR product may be subcloned and sequenced to ensure that the amplified sequences represent the sequences of a P-II gene. The PCR fragment may then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment may be labeled and used to screen a cDNA library, such as a plant cDNA library. Alternatively, the labeled fragment may be used to isolate genomic clones via the screening of a genomic library.

PCR technology may also be utilized to isolate full length cDNA sequences. For example, RNA may be isolated, following standard procedures, from an appropriate cellular or tissue source (i.e., one known, or suspected, to express the P-II gene. A reverse transcription reaction may be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNAase H, and second strand synthesis may then be primed with a poly-C primer. Thus, cDNA sequences upstream of the amplified fragment may easily be isolated. For a review of cloning strategies which may be used, see e.g., Sambrook et al., 1989, *supra*.

The P-II gene sequences may additionally be used to isolate mutant P-II gene alleles. Such mutant alleles may be isolated from plant species either known or proposed to have a genotype which contributes to the plant growth rate or nitrogen metabolism. Mutant alleles and mutant allele products may then be utilized in the therapeutic and diagnostic systems described below. Additionally, such P-II gene sequences can be used to detect P-II gene regulatory (e.g., promoter or promoter/enhancer) defects which can affect plant growth.

A cDNA of a mutant P-II gene may be isolated, for example, by using PCR, a technique which is well known to those of skill in the art. In this case, the first cDNA strand may be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known or suspected to be expressed in a plant species putatively carrying the mutant P-II allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal gene. Using these two primers, the product is then amplified via PCR, cloned into a suitable vector, and subjected to DNA sequence analysis through methods well known to those of skill in the art. By comparing the DNA sequence of the mutant P-II allele to that of the normal P-II allele, the mutation(s) responsible for the loss or alteration of function of the mutant P-II gene product can be ascertained.

Alternatively, a genomic library can be constructed using DNA obtained from a plant species suspected of or known to carry the mutant P-II allele, or a cDNA library can be constructed using RNA from a tissue known, or suspected, to express the mutant P-II allele. The normal P-II gene or any suitable fragment thereof may then be labeled and used as a probe to identify the corresponding mutant P-II allele in such libraries. Clones containing the mutant P-II gene sequences may then be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known, or suspected, to express a mutant P-II allele in a plant species suspected or known to carry such a mutant allele. In this manner, gene products made by the putatively mutant tissue may be expressed and screened

using standard antibody screening techniques in conjunction with antibodies raised against the normal P-II gene product, as described, below, in Section 5.3. (For screening techniques, see, for example, Harlow, E. and Lane, eds., 1988, "Antibodies: A Laboratory Manual", Cold Spring Harbor Press, Cold Spring Harbor.) Additionally, screening can be accomplished by screening the labeled P-II fusion proteins. It is noted where a P-II mutation results in an expressed gene product with altered function (e.g., as a result of a missense or a frameshift mutation), a polyclonal set of antibodies to P-II are likely to cross-react with the mutant P-II gene product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to sequence analysis according to methods well known to those of skill in the art.

The invention also encompasses nucleotide sequences that encode mutant P-II, peptide fragments of the P-II, truncated P-II, and P-II fusion proteins. These include, but are not limited to nucleotide sequences encoding mutant P-II described *infra*.

Due to the degeneracy of the nucleotide coding sequences, other DNA sequences which encode substantially the same amino acid sequences as depicted in FIG. 1 may be used in the practice of the present invention for the cloning and expression of P-II. Such alterations include deletions, additions or substitutions of different nucleotide residues resulting in a sequence that encodes the same or functionally equivalent gene product. The gene product may contain deletions, additions or substitutions of amino acid residues within the sequence, which result in a silent change thus producing a bioactive product. Such amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues involved. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; amino acids with uncharged polar head groups having similar hydrophilicity values include the following: leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine; phenylalanine, tyrosine.

The genomic sequences for P-II may be obtained from any plant cell source, whereas mRNA for preparation of cDNA copies may be obtained from cell sources that produce P-II. For example, parts of plants (e.g., leaves, stems, roots, nodules, cotyledons, seeds, fruits, etc.) may be ground and used as the source for extracting DNA or RNA. Preferably, the RNA is isolated from shoots, roots or flowers after exposure to light or treatment with sucrose. Alternatively, plant cell lines can be used as a convenient source of DNA or RNA. Genetically engineered microorganisms or cell lines containing P-II coding sequences, such as the deposited embodiments described herein, may be used as a convenient source of DNA for this purpose.

The P-II coding sequence may be obtained by cDNA cloning of RNA isolated and purified from such cellular sources or by genomic cloning. Either cDNA or genomic libraries may be prepared from the DNA fragments generated using techniques well known in the art, including but not limited to the use of restriction enzymes. The fragments which encode P-II may be identified by screening such libraries with a nucleotide probe that is substantially complementary to any portion of the nucleotide sequences depicted in FIGS. 12 and 13. Although portions of the coding sequence may be utilized, full length clones, i.e., those containing the entire coding region for P-II, may be preferable for expression. To these ends, techniques well known to those skilled in the art for the isolation of DNA,

generation of appropriate restriction fragments, construction of clones and libraries, and screening recombinants may be used. For a review of such techniques see, for example, Sambrook et al., 1989, *Molecular Cloning A Laboratory Manual* 2nd. ed., Cold Spring Harbor Press, N.Y. Alternatively, oligonucleotides derived from P-II sequences could be used as heterologous primers in PCR (polymerase chain reactions), to generate cDNA or genomic copies of P-II sequences from other species. For a review of such PCR techniques, see for example, Gelfand, D. H., 1989, "PCR Technology. Principles and Applications for DNA Amplification," Ed., H. A. Erlich, Stockton Press, N.Y.; and "Current-Protocols in Molecular Biology," Vol. 2, Ch. 15, Eds. Ausubel et al., John Wiley & Sons, 1988.

In an alternate embodiment of the invention, the coding sequences of FIGS. 12 and 13 could be synthesized in whole or in part, using chemical methods well known in the art. See, for example, Caruthers, et al., 1980, *Nuc. Acids Res. Symp. Ser.* 7:215-233; Crea and Horn, 180, *Nuc. Acids Res.* 9(10): 2331; Matteucci and Caruthers, 1980, *Tetrahedron Letters* 21:719; and Chow and Kempe, 1581, *Nuc. Acids Res.* 9(12) 2807-2817. Alternatively, the protein itself could be produced using chemical methods to synthesize the amino acid sequences depicted in FIGS. 12 and 13 in whole or in part. For example, peptides can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography. (E.g., see, Creighton, 1983, *Proteins Structures And Molecular Principles*, W. H. Freeman and Co., N.Y. pp. 50-60). The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, 1983, *Proteins, Structures and Molecular Principles*, W. H. Freeman and Co., N.Y., pp. 34-49).

In the specific embodiments described in the examples herein, the Arabidopsis P-II coding sequence was obtained by direct cloning of genomic sequences using a heterologous P-II cDNA probe from Ricinus. The P-II genomic sequence was used in turn to obtain an Arabidopsis P-II cDNA clone. Homology between P-II and bacterial PII was determined to be approximately 37% to 50%, suggesting that bacterial PII and (P-II) are evolutionarily related, yet quite distinct molecules (FIG. 1). Southern blot analysis of Arabidopsis nuclear DNA revealed that P-II is encoded by a single or low-copy number gene (FIG. 2). Northern blot analysis of Arabidopsis RNA revealed that the P-II is expressed in shoots, roots and flowers (FIG. 3). Our analysis of P-II mRNA has shown that P-II mRNA accumulates in these organs after induction by light or sucrose (FIGS. 4 and 5, respectively).

### 5.2. EXPRESSION OF P-II PROTEIN

The invention also encompasses (a) DNA vectors that contain any of the foregoing P-II coding sequences and/or their complements (i.e., antisense); (b) DNA expression vectors that contain any of the foregoing P-II coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences; and (c) genetically Engineered host cells that contain any of the foregoing P-II coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell. As used herein, regulatory elements include but are not limited to inducible and non-inducible promoters, enhancers, operators and other elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include but are not limited to the promoters derived from the genome of plant cells (e.g., heat shock promoters; the promoter for the

small subunit of RUBISCO; the promoter for the chlorophyll a/b binding protein) or from plant viruses (e.g., the 35S RNA promoter of CaMV; the coat protein promoter of tobacco mosaic virus (TMV), cytomegalovirus hCMV immediate early gene, the early or late promoters of SV40 adenovirus, the lac system, the trp system, the TAC system, the TRC system, the major operator and promoter regions of phage A, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase, the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

In order to express a biologically active P-II, the nucleotide sequence coding for P-II, or a functional equivalent as described in Section 5.1 supra, is inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. The P-II gene product as well as host cells, cell lines or plants transfected or transformed with recombinant P-II expression vectors can be used for a variety of purposes. These include but are not limited to generating antibodies (i.e. monoclonal or polyclonal) that define P-II; creating mutant P-II proteins which are constitutively active; creating transgenic plants containing such constitutively active mutant P-II genes; (creating mutant P-II proteins which are resistant to herbicides; creating transgenic plants containing such herbicide resistant mutant P-II genes); or creating transgenic plants which over-express P-II and demonstrate herbicide resistance; screening and selecting for P-II inhibitors or activators which may be used as herbicides.

### 5.3. CONSTRUCTION OF EXPRESSION VECTORS CONTAINING THE P-II CODING SEQUENCE

Methods which are well known to those skilled in the art can be used to construct expression vectors containing the P-II coding sequence and appropriate transcriptional/translational control signals. These methods include in vitro recombinant DNA techniques, synthetic techniques and in vivo recombination/genetic recombination. See, for example, the techniques described in Sambrook et al., 1989, *Molecular Cloning A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory, N.Y.).

A variety of host-expression vector systems may be utilized to express the P-II coding sequence. These include but are not limited to microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing the P-II coding sequence; yeast transformed with recombinant yeast expression vectors containing the P-II coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the P-II coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CAMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing the P-II coding sequence; or animal cell systems infected with recombinant virus expression vectors (e.g., adenovirus, vaccinia virus) containing the P-II coding sequence.

The expression elements of these vectors vary in their strength and specificities. Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used in the expression vector. For example, when cloning in bacterial systems, inducible promoters such as  $\pi$  of bacteriophage  $\lambda$ ,  $\text{plac}$ ,  $\text{ptrp}$ ,  $\text{ptac}$  ( $\text{ptrp-lac}$  hybrid promoter) and the like may be used; when cloning in insect cell systems, promoters such as the bacu-

lovirus polyhedrin promoter may be used; when cloning in plant cell systems, promoters derived from the genome of plant cells heat shock promoters; the promoter for the small subunit of RUBISCO; the promoter for the chlorophyll a/b binding protein) or from plant viruses (e.g., the 35S RNA promoter of CAMV; the coat protein promoter of TMV) may be used; when cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted P-II coding sequence.

In bacterial systems a number of expression vectors may be advantageously selected depending upon the use intended for the P-II expressed. For example, when large quantities of P-II are to be produced for the generation of P-II antibodies, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include but are not limited to the *E. coli* expression vector pUR278 (Ruther et al., 1983, *EMBO J.* 2:1791), in which the P-II coding sequence may be ligated into the vector in frame with the lac Z coding region so that a hybrid P-II-lac Z protein is produced; pIN vectors (Inouye & Inouye, 1985, *Nucleic acids Res.* 13:3101-3109; Van Heeke & Schuster, 1989, *J. Biol. Chem.* 264:5503-5509); and the like. However, where the P-II expression vector is to be used in an Asn-gln B- host for complementation assays described infra, the expression of unfused P-II using expression vectors with few or no host genotype requirements, including, but not limited to vectors such as ptac12, (Amann et al., 1983, *Gene* 25:167) and the like may be preferred.

In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review see, *Current Protocols in Molecular Biology*, Vol. 2, 1988, Ed. Ausubel et al., Greene Publish. Assoc. & Wiley Interscience, Ch-13; Grant et al., 1987, *Expression and Secretion Vectors for Yeast*, in *Methods in Enzymology*, Eds. Wu & Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp.516-544; Glover, 1986, *DNA Cloning*, Vol. II, IRL Press, Wash., D.C., Ch. 3; and Bitter, 1987, *Heterologous Gene Expression in Yeast*, *Methods in Enzymology*, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684; and *The Molecular Biology of the Yeast Saccharomyces*, 1982, Eds. Strathern et al., Cold Spring Harbor Press, Vols. I and II. For complementation assays in yeast, pea cDNAs for P-II may be cloned into yeast episomal plasmids (YEep) which replicate autonomously in yeast due to the presence of the yeast 2  $\mu$  circle. The P-II sequence may be cloned behind either a constitutive yeast promoter such as ADH or LEU2 or an inducible promoter such as GAL (*Cloning in Yeast*, Chpt. 3, R. Rothstein In: *DNA Cloning Vol.11, A Practical Approach*, Ed. DM Glover, 1986, IRL Press, Wash., D.C.). Constructs may contain the 5' and 3' non-translated regions of the cognate plant mRNA or those corresponding to a yeast gene. YEep plasmids transform at high efficiency and the plasmids are extremely stable. Alternatively vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

In cases where plant expression vectors are used, the expression of the P-II coding sequence may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson et al., 1984, *Nature* 310:511-514), or the coat protein promoter of TMV (Takamatsu et al., 1987, *EMBO J.* 6:307-311) may be used; alternatively, plant promoters such

as the small subunit of RUBISCO (Coruzzi et al., 1984, *EMBO J.* 3:1671-1680; Broglie et al., 1984, *Science* 224:838-843); or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B (Gurley et al., 1986, *Mol. Cell. Biol.* 6:559-565) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, etc. For reviews of such techniques see, for example, Weissbach & Weissbach, 1988, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp. 421-463; and Grierson & Corey, 1988, *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9.

An alternative expression system which could be used to express P-II is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The P-II coding sequence may be cloned into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the P-II coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (E.g., see Smith et al., 1983, *J. Virol.* 46:584; Smith, U.S. Pat. No. 4,215,051).

In cases where an adenovirus is used as an expression vector, the P-II coding sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing P-II in infected hosts. (E.g., See Logan & Shenk, 1984, *Proc. Natl. Acad. Sci. (USA)* 81:3655-3659). Alternatively, the vaccinia 7.5K promoter may be used. (E.g., see Mackett et al., 1982, *Proc. Natl. Acad. Sci. (USA)* 79:7415-7419; Mackett et al., 1984, *J. Virol.* 49:857-864; Panicali et al., 1982, *Proc. Natl. Acad. Sci.* 79:4927-4931).

Specific initiation signals may also be required for efficient translation of inserted P-II coding sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where the entire P-II gene, including its own initiation codon and adjacent sequences, are inserted into the appropriate expression vectors, no additional translational control signals may be needed. However, in cases where only a portion of the P-II coding sequence is inserted, exogenous translational control signals, including the ATG initiation codon, must be provided. Furthermore, the initiation codon, must be in phase with the reading frame of the P-II coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter et al., 1987, *Methods in Enzymol.* 153:516-544).

In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Expression driven by certain promoters can be elevated in the presence of certain inducers, (e.g., zinc and cadmium ions for metallothionein promoters). Therefore, expression of the genetically engineered P-II

may be controlled. This is important if the protein product of the cloned foreign gene is lethal to host cells. Furthermore, modifications (glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post translational processing and modification of proteins. Appropriate cells lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed.

One way to select for the expression of a biologically active P-II gene product, be it wild type, mutated or altered, and/or to screen for new herbicides, is to introduce an appropriate expression vector containing the P-II coding sequence into PII-minus strains of bacteria. The ability of an P-II cDNA to complement the host cells, indicates the expression and formation of an active P-II protein. A number of different PII-minus strains may be used for this purpose including but not limited to *E. coli*, *Klebsiella pneumoniae* and *Synechococcus* name but a few. Heterologous host systems may require the use of a second selectable marker as a means for selecting incorporation of the vector., e.g., in cases where P-II may not complement the PII-minus phenotype. To this end, dominant selectable markers such as antibiotic-resistance genes may be used.

#### 5.4. IDENTIFICATION OF TRANSFECTANTS OR TRANSFORMANTS EXPRESSING THE P-II GENE PRODUCT AND ISOLATION OF P-II

The host cells which contain the P-II coding sequence and which express the biologically active P-II gene product may be identified by these general approaches: (a) DNA-DNA hybridization; (b) the presence or absence of "marker" gene functions; (c) assessing the level of transcription as measured by the expression of P-II mRNA transcripts in the host cell; and (d) detection of the P-II gene product as measured by immunoassay or by its biological activity; or (e) phenotypic rescue.

In the first approach, the presence of the P-II coding sequence inserted in the expression vector can be detected by DNA-DNA hybridization using probes comprising nucleotide sequences that are homologous to the P-II coding sequences substantially as shown in FIGS. 12 and 13 or portions or derivatives thereof.

In the second approach, the recombinant expression vector/host system can be identified and selected based upon the presence or absence of certain "marker" gene functions (e.g., resistance to antibiotics, resistance to methotrexate, transformation phenotype, occlusion body formation in baculovirus, etc.). For example, if the P-II coding sequence is inserted within a marker gene sequence of the vector, recombinants containing the P-II coding sequence can be identified by the absence of the marker gene function. Alternatively, a marker gene can be placed in tandem with the P-II sequence under the control of the same or different promoter used to control the expression of the P-II coding sequence. Expression of the marker in response to induction or selection indicates expression of the P-II coding sequence. Two marker gene constructs which may be of particular-value for monitoring promoter activity in plant cells and plants are the bacterial glucuronidase gene, GUS (Jefferson et al., 1987, *EMBO J.* 6:3901-3908) or the luciferase gene (Ow et al., 1987, *Science*, 234:856-859).

In the third approach, transcriptional activity for the P-II coding region can be assessed by hybridization assays. For example, RNA can be isolated and analyzed by Northern blot using a probe homologous to the P-II coding sequence or particular portions thereof substantially as shown in

FIGS. 12 and 13. Alternatively, total nucleic acids of the host cell may be extracted and assayed for hybridization to such probes.

In the fourth approach, the expression of the P-II protein product can be assessed immunologically, for example by Western blots immunoassays such as radioimmuno-precipitation, enzyme-linked immunoassays and the like.

In the fifth approach, the production of a biologically active P-II gene product can be assessed by the complementation assay, in which a PII-minus bacterial transformed or transfected with the P-II expression vector will rescue the mutant phenotype. As previously explained, this may be used in conjunction with a separate selectable marker to ensure incorporation of the vector. In the sixth approach, mutant P-II which is constitutively active can be selected using the complementation assay expression system by exposing the clones which express mutant P-II to various concentrations of sucrose.

Once a clone that produces high levels of biologically active P-II is identified, the clone may be expanded and used for a variety of ends; e.g., production of P-II which may be purified using techniques well known in the art including but not limited to immunoaffinity purification, chromatographic methods including high performance liquid chromatography, and the like; screening herbicides; and engineering transgenic plants which have high nitrogen assimilatory efficiency.

#### 5.5. USES OF P-II GENE AND GENE PRODUCT

Our studies concerning P-II mRNA accumulation have highlighted the importance of the P-II mRNAs in plant nitrogen metabolism. The P-II cDNA clones can be used to characterize the P-II gene product; to express P-II so that antibodies which define P-II gene product can be produced; to screen and develop new herbicides; to develop herbicide resistant plants, nitrogen efficient plants.

#### 5.6. PRODUCTION OF ANTIBODIES THAT DEFINE P-II

Expressed gene products may be used to produce antibodies that define P-II. These antibodies can be used in fractionation studies to define the cellular and tissue site of action of P-II polypeptide. For full length cDNA clones, the corresponding proteins may be produced in *E. coli* in sufficient quantities to allow characterization in terms of substrate preference (i.e. sucrose and other sugars), Km, sensitivity to inhibitors, etc.

Such antibodies may be produced by any method known in the art, including, but not limited to injection of P-II into mice, rats, rabbits, or other host species for production of polyclonal antisera. Various adjuvants may be used to increase the immune response. These include, but are not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, and oil emulsions. For a review of such techniques, see Harlow & Lane, 1988, *Antibodies A Laboratory Manual*, Cold Spring Harbor Laboratory.

Monoclonal antibodies, or fragments thereof, can be prepared by any technique which provides for the production of antibody molecules by continuous cell lines in culture. Such techniques include but are not limited to the hybridoma technique first developed by Kohler and Millstein (1975, *Nature* 256:495-497), the human B-cell hybridoma technique (Kozbor et al., 1983, *Immunology Today* 4:72), and the EBV-hybridoma technique for production of human monoclonal antibodies (Cole et al., 1985, *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96).

Antibody fragments which contain the idiotype of the molecule could be generated by known techniques. For example, such fragments include but are not limited to: the F(ab')<sub>2</sub> fragment which can be produced by pepsin digestion of the antibody molecule; the Fab' fragments which can be generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragment, and the 2 Fab or Fab fragments which can be generated by treating the antibody molecule with pepsin and a reducing agent.

#### 5.7. DEVELOPMENT OF NEW HERBICIDES

P-II cDNA clones can be used in functional complementation assays described in Section 5.1, supra, designed to screen for new herbicides as follows. Full length P-II cDNAs cloned in the appropriate expression vectors can be introduced into P-II-minus strains of bacteria as previously described. The ability of a P-II cDNA clone to rescue the P-II-minus mutant cells would indicate that the cDNA encodes an P-II polypeptide which is active in these heterologous environments.

The P-II cDNAs expressed in P-II-minus mutants of bacteria described previously provide an in vivo system to select for herbicides which selectively act on P-II. To this end, known concentrations of a test substance can be added to the growth media in order to select those substances which inhibit cell growth as an indication of P-II inhibitory activity.

#### 5.8. DEVELOPMENT OF HERBICIDE RESISTANT PLANTS AND NITROGEN-EFFICIENT PLANTS

The in vivo expression system described above could be modified and used to select for mutations in the P-II structural gene which are constitutively active or confer growth resistance to the new herbicides. Mutations may be introduced into the P-II cDNA in vivo or in vitro using techniques well known in the art, including, but not limited to, radiation, chemical mutation, site-specific mutations, etc. For example, see Tilghman and Levine, 1987, in *Gene Transfer*, Kucherlaputi, Ed., Plenum Pub., N.Y., pp. 189-221). Expression vectors containing such altered or mutated P-II coding sequences can be used in P-II-minus bacterial hosts in the complementation assay described above to identify clones containing coding sequences that specify and express biologically active, altered P-II. Clones which produce constitutively active or herbicide resistant P-II can be selected by growth in the absence of inducer or the presence of the herbicide, respectively. In this way, clones which produce mutant P-II can be screened for constitutive activity or resistance to new herbicides.

Plants may then be engineered for enhanced nitrogen utilizations or resistance to herbicides using appropriate constructs containing the mutated P-II genes which over-express wild-type P-II or antisense to P-II which results in down regulation of P-II expression. In either case, such transgenic plants may be constructed using methods well known to those skilled in the art including but not limited to techniques involving the use of the Ti plasmids (eg., *Agrobacterium rhizogenes*), plant viruses, electroporation, direct transformation, microinjection, etc. By way of example, and not by way of limitation, the binary Ti vector system could readily be used to this end (Bevan, 1984, *Nuc. Acids Res.* 12:8711-8721). This vector contains a selectable marker, neomycin phosphotransferase (kanR), under direction of the nopaline synthetase promoter (nos) for KanR selection in plants, and unique cloning sites for EcoRI, HindIII, BamHI, SmaI, SalI, KpnI, baI, SstI. Using this system, DNA fragments cloned into one of the unique cloning sites located

*faciens* LB4404 or GU3010 harboring a resident disarmed Ti plasmid which will provide vir gene required for T-DNA transfer in trans. Another vector system which could be used is the pMON505 intermediate binary Ti transformation vector (Horsch & Klee, 1986, *Proc. Natl. Acad. Sci.* 83, 4428-4432; Horsch et al., 1985 *Science*, 227:1229-1231). For reviews of such techniques see, for example, Weissbach & Weissbach, 1988, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp. 421-463; and Grierson & Corey, 1988, *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9. P-II genes altered in vivo or in vitro to overproduce wild type P-II or constitutively active or herbicide resistant forms of P-II enzymes may be used as dominant selectable markers for transformation systems which include organisms such as bacteria, algae, yeast, neurospora and plants.

By way of illustration, enhanced nitrogen utilization or herbicide resistant plants can be engineered in at least two ways: P-II cDNAs which contain mutations conferring constitutive activity or herbicide resistance as measured in the screening assay above, can be introduced into transgenic plants under the transcriptional regulation of a strong constitutive promoter (e.g., 35S promoter of CaMV) or an inducible promoter (e.g., the promoter of the small subunit of RUBISCO; a heat shock promoter, etc.). Mutant P-II produced in the transgenic plants will confer enhanced nitrogen utilization or herbicide resistance. Alternatively, enhanced nitrogen utilization or herbicide resistant plants could be developed by the over-expression or underexpression of the wild-type P-II cDNA. The overexpression or antisense of P-II in these transgenic plants will produce plants with enhanced nitrogen utilization properties or resistant to high levels of a herbicide specific of P-II.

#### 5.9. THE P-II PROMOTER

As described above and exemplified below, the P-II promoter directs expression in shoots, roots and flowers. This promoter can be used to direct the regulated expression of heterologous gene sequences in appropriate hosts. The P-II promoter described herein is not only inducible by light or sucrose, but drives high levels of transcription in flowers.

The P-II promoter may be isolated from the P-II genomic clones.

The organ-specific and photo inducible P-II promoter can be used in a variety of DNA vectors to drive the expression of heterologous sequences ligated after the transcription start site. Heterologous sequences ligated downstream of the transcription start site, containing a portion of the leader sequence, will require translational control elements such as the ATG start signal, and ribosome binding sites etc. optionally, the heterologous sequences may also be ligated in a translational fusion to marker genes such  $\beta$ -galactosidase, GUS, luciferase, etc. to produce fusion proteins.

Such expression vectors may be used in plant cell culture or in transgenic plants to direct high level expression of the heterologous sequence in an inducible, temporal, or organ-specific fashion.

## 6. EXAMPLE

### ARABIDOPSIS AND CASTOR BEAN P-II GENE

A plant gene GLB1 encoding a PII homologue was identified, the first time a PII gene has been identified in a eukaryote. This plant PII-homologue was identified as a cDNA present in an Expressed Sequence Tag (EST) library

of castor (*Ricinus communis*) (van de Loo et al., 1995, Plant Physiol.) and was subsequently cloned from Arabidopsis. In this example, the two plant GLB1 cDNAs encoding PII are characterized and it is shown that their deduced protein sequences have striking identity to microbial PII. It is also demonstrated that the regulation of GLB1 mRNA levels by light and by metabolites correlates with the activation of putative downstream genes such as chloroplastic GS2 (GLN2) (Faure et al., 1994, *Plant J.* 5:481–491; Edwards, J. W. and Coruzzi, G. M., 1989, *Plant Cell* 1:241–248). Combined with previous findings that plant nitrogen assimilatory genes are regulated by carbon and nitrogen metabolites (Faure et al., 1994, *Plant J.* 5:481–491; Lam et al., 1994, *Plant Physiol.* 106:1347–135.; Vincentz et al., 1993, *Plant J.* 3:315–324), these results suggest that PII may be the first nitrogen regulatory protein identified in plants controlling the assimilation of inorganic nitrogen into organic form. As nitrogen assimilation into organic form may be a critical rate-limiting element in plant growth, the discovery of a gene that regulates this process may have important implications to improving nitrogen-use efficiency in plants.

#### MATERIALS AND METHODS

##### Plant Materials

The plant tissues used in all experiments were from *Arabidopsis thaliana* Columbia ecotype. Plants used for genomic DNA Extraction, RNA isolation from different organs were grown in soil. Others were grown in a semi-hydroponic system developed by Applicants (I. Oliveira and G. Coruzzi, unpublished). This system contains MS salt mixture (Sigma) plus 0.4% agar in Plantcons (ICN Laboratories Inc.). Arabidopsis seeds were sown on Nylon grids (250  $\mu$ M) (Tetko Inc.) placed on the surface of the media. Plants were grown in EGC growth chambers at 45  $\mu$ E m<sup>-2</sup> sec<sup>-1</sup> on a 16 h light/8 h dark cycle, unless otherwise noted. Isolation of Castor and Arabidopsis GLB1 cDNA Clones

A  $\lambda$ ZAPII library containing cDNAs from developing endosperm and embryos of castor (*Ricinus communis* L.) seeds was used for mass-sequencing (van de Loo et al., 1995, *Plant Physiol.*). Out of 743 clones sequenced, one clone showed sequence similarity to nitrogen-regulatory protein PII of *Synechococcus* and other bacteria (Tsinoremas et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:4565–4569). This 0.85 kb cDNA called pRc-GLB1 was used to make a <sup>32</sup>P-dCTP labeled probe by random priming and used to screen an Arabidopsis genomic library at a reduced stringency as follows. The hybridization was performed in QuickHyb (Stratagene) at 50° C. for 1 hr. Filters were washed with 2 $\times$ SSC, 0.1% SDS, 5 min twice at room temperature followed by 0.5 $\times$ SSC, 0.1% SDS wash at 50° C. for 30 min. One Arabidopsis genomic GLB1 clone with a 3 kb DNA insert was isolated. A 1.1 kb NcoI-NcoI DNA fragment of the genomic clone which strongly hybridized to the castor GLB1 cDNA was used to screen an Arabidopsis silique cDNA library. Under high-stringency hybridization conditions, two Arabidopsis cDNA clones were obtained from 2 $\times$ 10 plaques. These Arabidopsis GLB1 cDNAs were sequenced using the Sequenase method (U.S. Biochemical Corp.). DNA sequences were analyzed by the GCG Sequence Analysis software package (Genetics Computer Group Inc., Madison, Wis.). 5' RACE was performed using Gibco-BRL 5' RACE system. Reverse transcription was performed with 1  $\mu$ g total RNA from light-grown plants. Nested primers for RACE were MH7 (GCAAGATGGTCCGGGAATGTC) (SEQ ID NO:17), MH8 (CGACAGGTAAAACACGACTG) (SEQ ID NO:18) and MH9 (GGTCTGACAAATTGCTTCCAC) (SEQ ID NO:19). PCR conditions were 94° C. 2 min, followed by 30 cycles

(94° C. 20 sec, 55° C. 30 sec, and 72° C. 40 sec). PCR products were subcloned using the pCR-Script kit of Stratagene.

##### Analyses of DNA and RNA

Arabidopsis genomic DNA was isolated according to the procedure described by Ausubel et. al, 1987, In: *Current Protocols in Molecular Biology*, (Greene Publishing Assoc. & John Wiley & Son), except that the buffer used was 8.75 M urea, 438 mM NaCl, 62.5 mM Tris-HCl pH 8.0, 62.5 mM EDTA. RNA was isolated using a phenol extraction protocol (Jackson, A. O. and Larkins, B. A., 1976, *Plant Physiol.* 57:5–10). Southern blot and Northern blot were performed as described (Sambrook et al., *Molecular cloning: a laboratory manual*, 3rd ed. (Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory)). For detection of GLB1 mRNA, membranes were hybridized with single-stranded DIG-labeled probes made from pAt-GLB1 cDNA using PCR (Myerson, D., 1991, *Biotechniques* 10:35–38). To generate probes approximately 540 nucleotides in length covering the PII coding region and partial 3' non-coding region, MH5 (GAAACCAAACACAGACTCC) (SEQ ID NO:20) and MH6 (CCGAGTAATAACAGTCGTC) (SEQ ID NO:21) primers were used. The control DIG-labeled probes for GLN2, P-PIIN1 and 18 rRNA were made by random priming (Boehringer Mannheim).

##### Computer Analysis

Geneworks (Intelligenetics Inc) was used to align the polypeptide sequences of plant and bacterial PII's. Phylogenetic analysis was performed using the heuristic option of PAUP 3.1.1 (phylogenetic analysis using parsimony) (Stewart, C-B., 1993, *Nature* 361:603–607). Archaeobacterial PII-like protein 1 was selected as the outgroup. To generate the appropriate alignment for use in PAUP, insertions and deletions were introduced using 'MALIGNED' (Clark, S. P., 1992, *Comput. Appl. Biosci.* 8:535–538). The PII sequences for bacteria and archaeobacteria were obtained from GenBank.

#### RESULTS

##### Cloning and Identification of Arabidopsis and Castor PII-homologue cDNAs

By mass-sequencing of portions of cDNAs from a castor developing seed library, a 0.85 kb cDNA clone, pCRS852, was identified which showed significant similarity to the nitrogen-regulatory protein PII of enteric bacteria encoded by *glnB*. This plant cDNA clone was named pRc-GLB1 and was further analyzed and the sequence of the entire clone determined. The castor *glnB*-like cDNA was used to isolate an Arabidopsis cDNA homologue. Two independent Arabidopsis clones with same size cDNA insert, 0.82 kb, were obtained. The nucleotide sequence of these two Arabidopsis cDNA clones encoding P-II revealed that they possess identical sequences in the coding region and non-coding regions. One clone extends further at the 3' end and appears to use a different polyadenylation site (data not shown). Thus, these two Arabidopsis cDNAs were considered to be derived from the same GLB1 gene. These Arabidopsis cDNAs encoding P-II are judged to be full-length for the following reasons. First, the size of the cDNA is consistent with that predicted for the mRNA based on Northern blot analysis. Second, three independent 5'-RACE products all showed an in-frame TAA stop codon 45 nucleotides upstream of the first ATG (data not shown).

The Arabidopsis PII (GLB1) gene maps to the top arm of chromosome 4. RFLP and recombinant inbred lines of Arabidopsis were used to map the PII (GLB1) gene to the top arm of chromosome 4 at position 10.8 (FIG. 10). Plant PII has Significant Similarity to Microbial PII Proteins.



The P-PII polypeptide encoded by the Arabidopsis GLB1 cDNA is 196 amino acids long with a molecular weight of 22 kD daltons. The region of homology to bacterial PII occurs between amino acids 74 and 186 of the Arabidopsis P-II protein. This region encompasses the entire microbial PII protein (FIG. 1A). In this region, the overall identity between plant and *E. coli* PII is 50.9%. The identity between P-II and *E. coli* PII is extremely high in two "signature domains" (Sor., H. S. and Rhee, S. G., 1987, J. Biol. Chem. 262:8690-8695) conserved amongst all PII proteins (85%) (FIG. 1A). Within the region corresponding to *E. coli* PII (residues 74-186 of Arabidopsis P-II) the two plant cDNAs are 90.3% identical (FIG. 1A). Beyond this region, the identity between these two P-II proteins decreases to 32.9% at the amino termini and 40% at the carboxy-termini (data not shown). When the deduced P-II amino acid sequences are compared with those of microbial glnB genes encoding PII, high overall identities are found with *E. coli* (50.9%) (Son, H. S. and Rhee, S. G., 1987, J. Biol. Chem. 262:8690-8695), *Synechococcus* sp. PCC 7942 (56.4%) (Tsinoremas et al., 1991, Proc. Natl. Acad. Sci. USA 88:4565-4569), *Klebsiella pneumoniae* (50.9%) (Holtel, A. and Merrick, M., 1988, Mol. Gen. Genet. 215:134-138), *Azospirillum brasilense* (49.1%) (de Zamaroczy et al., 1990, Mol. Gen. Genet. 224:421-430), *Rhodobacter capsulatus* (48.2%) (Kranz et al. 1990, J. Bacteriol. 1;2:53-62. *Rhizobium leguminosarum* (47.7%) Colonna-Romano et al., 1987, Nucleic Acids Res. 15:1951-1964, *Bradyrhizobium japonicum* (47.7%) (Martin et al., 1989, J. Bacteriol. 171:5638-5645), *Bacillus subtilis* (38.6%) (Wray et al., 1994, J. Bacteriol. 176:108-114) and with *Methanococcus thermolithotrophicus* (archaeobacteria) glnB-like protein 1 (34.9%) (Souillard, N. and Sibold, L., 1989, Mol. Microbiol. 3:541-551).

In addition to the high overall identity amongst various PII proteins of bacteria and plants, the P-II proteins share extremely high regional identities over signature domains 1 and 2 (85%). Signature domain 1 contains the Tyr 51 residue which is post-translationally modified by uridylylation in enteric bacteria (Magasanik, B., 1988, TIBS 13:475-479). As the glnB mRNA is constitutively expressed in *E. coli* (van Heeswijk et al., 1993, Mol. Microbiol. 9:443-457). The activity of the PII protein product is regulated post-translationally by uridylylation. Although the uridylylation of Tyr51 is considered an important feature of signature domain 1 of bacterial PII proteins, exceptions do exist. In *Bacillus*, the PII-like protein does not contain this conserved tyrosine residue. However it is not known if this is a functional PII protein (Wray et al., 1994, J. Bacteriol. 176:108-114). In cyanobacteria, although PII possesses Tyr 51, its activity is regulated by phosphorylation at a serine residue (Forchhammer, K. and De Marsac, N. T., 1994, J. Bacteriol. 176:84-91). In both plant PII proteins, the Tyr 51 is replaced by a phenylalanine (FIG. 1). Interestingly, there is only one nucleotide change between tyrosine and phenylalanine codons. The lack of Tyr 51 in the P-II proteins suggests that the P-II protein may not be regulated post-translationally by uridylylation. The conserved signature domain-2 common to all bacterial PIIs, is extremely well conserved with Arabidopsis and castor PII proteins (FIG. 1A). The extremely high conservation between prokaryotic and eukaryotic PII suggests that this second domain plays an important, but as yet unknown role in PII function.

A phylogenetic analysis of the evolutionary relationship among different PII proteins in plants and bacteria was performed using a parsimony method (PAUP) (Stewart, C.-B., 1993, Nature 361:603-607). (FIG. 1B). Archaeobacteria

was used as the outgroup in this tree. The overall homology between plant PII protein and bacterial PII (50%) is significantly higher than plant PII is to the archaeobacteria (35%). However, the Tyr51 residue, which is conserved in the PII proteins of cyanobacteria and most of the eubacteria, is replaced by a Phe in both plants and archaeobacteria.

GLB1 Encoding a PII Homologue is a Single- or Low-copy Number Gene in Arabidopsis.

Arabidopsis genomic DNA was digested with several different restriction enzymes and probed with a DNA fragment from the GLB1 cDNA (FIG. 2). In each lane, one or several Arabidopsis genomic DNA fragments hybridized to the GLB1 cDNA clone. Preliminary restriction enzyme analysis of an Arabidopsis GLB1 genomic clone encoding PII was also performed. These data indicate that PII is truly encoded by a plant gene, GLB1, and that in Arabidopsis is a single- or low-copy number gene.

## 7. EXAMPLE

### INDUCTION OF PII mRNA IN RESPONSE TO LIGHT AND SUCROSE

These studies demonstrate that P-II mRNA is expressed in both photosynthetic and non-green Arabidopsis organs and that the expression of the P-II gene is induced by light and sucrose.

#### 7.1. THE GLB1 mRNA ENCODING PII IS EXPRESSED IN PHOTOSYNTHETIC AND NON-GREEN ARABIDOPSIS ORGANS

The expression of GLB1 mRNA in different organs of Arabidopsis was examined by Northern blot analysis. Arabidopsis GLB1 mRNA is detected in shoots, roots, and flowers (FIG. 3 lanes 1-3). The expression of GLB1 mRNA in different organs may reflect the function of the PII protein in each tissue. For example, primary ammonium assimilation via GS/GOGAT occurs primarily in photosynthetic organs. It is therefore reasonable to detect higher levels of GLB1 mRNA in leaves compared to roots (FIG. 3, lanes 1 & 2). The high level expression of GLB1 mRNA in developing flowers (FIG. 3, lane 3) and siliques with developing seeds (not shown) is consistent with the highly active nitrogen metabolism in these organs.

#### 7.2. ARABIDOPSIS GLB1 mRNA LEVELS ARE INDUCED BY LIGHT

Light appears to coordinate the expression of nitrogen assimilatory genes such as NR, GLN2, and GLU1 (encoding Fd-GOGAT) with that of photosynthetic genes such as rbcS. It was examined whether light also induced the accumulation of Arabidopsis GLB1 mRNA, a putative nitrogen-regulatory gene (FIG. 4A and GLB1 mRNA was shown to accumulate preferentially in light-grown plants compared to plants dark-adapted for 48 hrs (FIG. 4A, lanes 1 & 2). A time-course of light-induction was conducted and the identical Northern blot was probed simultaneously with DNA probes for GLB1 and its putative downstream target gene GLN2 (FIG. 4B). The induction of GLB1 and GLN2 mRNA levels are coordinately induced by light to maximal induced levels (FIG. 4B, lane 4). This finding is consistent with the hypothesis that the PII protein encoded by GLB1 may regulate downstream nitrogen assimilatory genes such as GLN2.

#### 7.3. LEVELS OF ARABIDOPSIS GLB1 MRNA ARE INDUCED BY SUCROSE

Light via phytochrome has been shown to have a direct effect on the expression of genes for nitrogen assimilatory enzymes such as nitrate reductase (Crawford et al., 1986, Proc. Natl. Acad. Sci. USA. 83:8073-8076), glutamine syn-

thetase (McGrath, R. B. and Coruzzi, G. M., 1991, *Plant J.* 1:275–280), and glutamate synthase, (Sakakibara, H. et al., 1991, *J. Biol. Chem.* 266:2028–2035). It appears that light-induced changes in carbon metabolites can also affect gene induction, as sucrose has been shown to induce the expression of NR (Vincentz et al., 1993, *Plant J.* 3:315–324), GLN2 (Faure et al., 1994, *Plant J.* 5:481–491) and GLU1, independent of light. Metabolic gene induction by sucrose links the expression of nitrogen assimilatory genes with the availability of carbon skeletons. As GLB1 mRNA levels are induced by light, we examined whether sucrose could also affect the expression of Arabidopsis GLB1 mRNA. Steady state levels of GLB1 mRNA are low in dark-adapted plants and are induced by light 10-fold (FIG. 5A, compare lanes 1 & 4). The effect of light on levels of GLB1 mRNA can be mimicked by the addition of 3% sucrose (FIG. 5A, lane 2), but not by mannitol, a non-metabolizable carbon source (FIG. 5A, lane 3). The effects of sucrose on the levels of GLB1 mRNA in dark-adapted Arabidopsis parallels the accumulation of mRNA for Arabidopsis chloroplast GS2 (GLN2) (Faure et al., 1994, *Plant J.* 5:481–491). and Fd-GOGAT genes (GLU1). Interestingly, sucrose can super-induce GLB1 mRNA accumulation beyond light-induced levels (FIG. 5A, lane 5). Again this effect is not obtained with mannitol (FIG. 5A, lane 6). Table 1 details the quantitation of GLB1 mRNA from two separate experiments. As sucrose acts independent of light to induce GLB1 mRNA, these results prove that the GLB1 gene is subjected to metabolic control.

Induction of PII (GLB1) mRNA by light/sucrose mirrors that of the putative downstream target nitrogen assimilatory genes GS2 (GLN2) and Fd-GOGAT (GLU1). PII mRNA is low in dark-adapted plants and is induced by sucrose in the absence of light, but not by mannitol, a non-metabolizable sugar (FIG. 6). Light induces PII mRNA and light plus sucrose super-induces PII mRNA (lane 5). Mannitol cannot superinduce PII mRNA above light levels. Nitrogen assimilatory genes GS2 and Fd-GOGAT GLU1, also show induction by sucrose or light, but are not super-induced by light and sucrose (FIG. 6).

PII induction by light and/or high sucrose, mirrors the induction of downstream nitrogen assimilatory genes for glutamine synthetase (GLN2), Fd-dependent glutamate synthase (Fd-GOGAT, GLU2), and aspartate aminotransferase (ASP2). By contrast, genes for asparagine synthetase (ASN1) and glutamate dehydrogenase are repressed by high sucrose (FIG. 7).

TABLE 1

Fold-induction of GLB1 mRNA levels induced by light and/or sucrose		
Lane	Treatments	Fold-Induction*
1.	Dark	0% Sucrose
1.0		
2.	Dark	3% Sucrose

TABLE 1-continued

Fold-induction of GLB1 mRNA levels induced by light and/or sucrose		
Lane	Treatments	Fold-Induction*
7.4		
4.	Light	0% Sucrose
10.2		
5.	Light	3% Sucrose
21.8		

\*GLB1 mRNA signals depicted in FIG. 5 plus a replicate experiment were quantified by densitometry. The average fold-induction relative to lane 1, FIG. 5, is listed.

## 8 EXAMPLE

## PII (GLB1) EXPRESSION IS REPRESSED BY HIGH ORGANIC NITROGEN, RELATIVE TO CARBON

This study demonstrates that the regulation of PII mirrors the regulation of the putative downstream nitrogen assimilatory gene for glutamine synthetase (GLN2) (FIG. 8). PII and its putative downstream nitrogen assimilatory genes (GLN2, GLU1 and ASP2) are induced by sucrose (FIG. 9, lane 1). Sucrose induction is repressed by the addition of organic nitrogen as asparagine, glutamine and glutamate (FIG. 9). By contrast, ASN1 and GDH1 are induced by these conditions. These results are consistent with a role for PII in the carbon:nitrogen sensing/signalling mechanism leading to changes in the expression of genes involved in nitrogen assimilation.

## 9. EXAMPLE

## TRANSGENIC TOBACCO PLANTS EXPRESSING ANTISENSE PII mRNA

Antisense Arabidopsis PII was introduced into tobacco by Agrobacterium-mediated DNA transfer. Shown are four independent transgenic tobacco lines which express high levels of PII antisense RNA (FIG. 11), lines 1, 2, 5 and 10). Phenotypic analysis of plants is ongoing. In addition, both sense and antisense PII Arabidopsis transgenic plants have been created in order to test whether the C:N sensing mechanism is perturbed in these transgenic plants.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entireties.

The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 21

-continued

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Phe Tyr Lys Val Glu Ala Ile Val Arg Pro Trp Arg Ile Gln Gln Val  
 1 5 10 15

Ser Ser Ala Leu Leu Lys Ile Gly Ile Arg Gly Val Thr Val Ser Asp  
 20 25 30

Val Arg Gly Phe Gly Ala Gln Gly Gly Ser Thr Glu Arg His Gly Gly  
 35 40 45

Ser Glu Phe Ser Glu Asp Lys Phe Val Ala Lys Val Lys Met Glu Ile  
 50 55 60

Val Val Lys Lys Asp Gln Val Glu Ser Val Ile Asn Thr Ile Ile Glu  
 65 70 75 80

Gly Ala Arg Thr Gly Glu Ile Gly Asp Gly Lys Ile Phe Val Leu Pro  
 85 90 95

Val Ser Asp Val Ile Arg Val Arg Thr Gly Glu Arg Gly Glu Lys Ala  
 100 105 110

Glu

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 113 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Phe Tyr Lys Val Glu Ala Ile Leu Arg Pro Trp Arg Val Ser Gln Val  
 1 5 10 15

Ser Ser Ala Leu Leu Lys Ile Gly Ile Arg Gly Val Thr Val Ser Asp  
 20 25 30

Val Arg Gly Phe Gly Ala Gln Gly Gly Ser Thr Glu Arg Gln Gly Gly  
 35 40 45

Ser Glu Phe Ser Glu Asp Lys Phe Val Ala Lys Val Lys Met Glu Ile  
 50 55 60

Val Val Ser Lys Asp Gln Val Glu Asp Val Ile Glu Lys Ile Ile Glu  
 65 70 75 80

Glu Ala Arg Thr Gly Glu Ile Gly Asp Gly Lys Ile Phe Leu Leu Pro  
 85 90 95

Val Ser Asp Val Ile Arg Val Arg Thr Gly Glu Arg Gly Asp Lys Ala  
 100 105 110

Glu

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 amino acids
- (B) TYPE: amino acid

-continued

(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Met Lys Lys Ile Asp Ala Ile Ile Lys Pro Phe Lys Leu Asp Asp Val
 1           5           10           15
Arg Glu Ala Leu Ala Glu Val Gly Ile Thr Gly Met Thr Val Thr Glu
           20           25           30
Val Lys Gly Phe Gly Arg Gln Lys Gly His Thr Glu Leu Tyr Arg Gly
           35           40           45
Ala Glu Tyr Met Val Asp Phe Leu Pro Lys Val Lys Ile Glu Ile Val
           50           55           60
Val Thr Asp Asp Ile Val Asp Thr Cys Val Asp Thr Ile Ile Arg Thr
           65           70           75           80
Ala Gln Thr Gly Lys Ile Gly Asp Gly Lys Ile Phe Val Phe Asp Val
           85           90           95
Ala Arg Val Ile Arg Ile Arg Thr Gly Glu Glu Asp Asp Ala Ala Ile
           100           105           110

```

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 112 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

```

Met Lys Lys Ile Asp Ala Ile Ile Lys Pro Phe Lys Leu Asp Asp Val
 1           5           10           15
Arg Glu Ala Leu Ala Glu Val Gly Ile Thr Gly Met Thr Val Thr Glu
           20           25           30
Val Lys Gly Phe Gly Arg Gln Lys Gly His Thr Glu Leu Tyr Arg Gly
           35           40           45
Ala Glu Tyr Met Val Asp Phe Leu Pro Lys Val Lys Ile Glu Ile Val
           50           55           60
Val Pro Asp Asp Ile Val Asp Thr Cys Val Asp Thr Ile Ile Arg Thr
           65           70           75           80
Ala Gln Thr Gly Lys Ile Gly Asp Gly Lys Ile Phe Val Phe Asp Val
           85           90           95
Ala Arg Val Ile Arg Ile Arg Thr Gly Glu Glu Asp Asp Ala Ala Ile
           100           105           110

```

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 111 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

Met Lys Lys Ile Glu Ala Ile Ile Lys Pro Phe Lys Leu Asp Glu Val
 1           5           10           15
Arg Ser Pro Ser Gly Val Gly Leu Gln Gly Ile Thr Val Thr Glu Ala

```

-continued

	20		25		30														
Lys	Gly	Phe	Gly	Arg	Gln	Lys	Gly	His	Thr	Glu	Leu	Tyr	Arg	Gly	Ala				
	35						40					45							
Glu	Tyr	Val	Val	Asp	Phe	Leu	Pro	Lys	Val	Lys	Val	Glu	Val	Val	Leu				
	50					55					60								
Ala	Asp	Glu	Asn	Ala	Glu	Ala	Val	Ile	Glu	Ala	Ile	Arg	Lys	Ala	Ala				
65					70				75					80					
Gln	Thr	Gly	Arg	Ile	Gly	Asp	Gly	Lys	Ile	Phe	Val	Ser	Asn	Val	Glu				
			85						90					95					
Glu	Val	Ile	Arg	Ile	Arg	Thr	Gly	Glu	Thr	Gly	Ile	Asp	Ala	Ile					
			100					105					110						

## (2) INFORMATION FOR SEQ ID NO:6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 111 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Lys	Lys	Ile	Glu	Ala	Ile	Ile	Lys	Pro	Phe	Lys	Leu	Asp	Glu	Val				
1			5					10					15						
Arg	Ser	Leu	Ser	Gly	Val	Gly	Leu	Gln	Gly	Ile	Thr	Val	Thr	Glu	Ala				
		20					25						30						
Lys	Gly	Phe	Gly	Arg	Gln	Lys	Gly	His	Thr	Asp	Leu	Tyr	Arg	Gly	Ala				
		35					40					45							
Glu	Tyr	Ile	Val	Asp	Phe	Leu	Pro	Lys	Val	Lys	Ile	Glu	Ile	Val	Ile				
	50					55				60									
Gly	Asp	Asp	Leu	Val	Glu	Arg	Ala	Ile	Asp	Ala	Ile	Arg	Arg	Ala	Ala				
65					70				75					80					
Gln	Thr	Gly	Arg	Ile	Gly	Asp	Gly	Lys	Ile	Phe	Val	Ser	Asn	Ile	Glu				
			85						90					95					
Glu	Ala	Ile	Arg	Ile	Arg	Thr	Gly	Glu	Ser	Gly	Leu	Asp	Ala	Ile					
			100					105					110						

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Lys	Lys	Ile	Glu	Ala	Ile	Ile	Lys	Pro	Phe	Lys	Leu	Asp	Glu	Val				
1			5					10					15						
Lys	Glu	Ala	Leu	His	Glu	Val	Gly	Ile	Lys	Gly	Ile	Thr	Val	Thr	Glu				
		20					25						30						
Ala	Lys	Gly	Phe	Gly	Arg	Gln	Lys	Gly	His	Thr	Glu	Leu	Tyr	Arg	Gly				
		35					40					45							
Ala	Glu	Tyr	Val	Val	Asp	Phe	Leu	Pro	Lys	Val	Lys	Ile	Glu	Val	Val				
	50					55					60								
Met	Glu	Asp	Ser	Leu	Val	Glu	Arg	Ala	Ile	Glu	Ala	Ile	Gln	Gln	Ala				
65					70				75					80					

-continued

Ala His Thr Gly Arg Ile Gly Asp Gly Lys Ile Phe Val Thr Pro Val  
85 90 95

Glu Glu Val Val Arg Ile Arg Thr Gly Glu Lys Gly Gly Asp Ala Ile  
100 105 110

## (2) INFORMATION FOR SEQ ID NO:8:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Lys Lys Val Glu Ala Ile Ile Lys Pro Phe Lys Leu Asp Glu Val  
1 5 10 15

Lys Glu Ala Leu Gln Glu Ala Gly Ile Gln Gly Leu Ser Val Ile Glu  
20 25 30

Val Lys Gly Phe Gly Arg Gln Lys Gly His Thr Glu Leu Tyr Arg Gly  
35 40 45

Ala Glu Tyr Val Val Asp Phe Leu Pro Lys Val Lys Ile Glu Met Val  
50 55 60

Leu Pro Asp Glu Met Val Asp Ile Ala Ile Glu Ala Ile Val Gly Ala  
65 70 75 80

Ala Arg Thr Glu Lys Ile Gly Asp Gly Lys Ile Phe Val Ser Ser Ile  
85 90 95

Glu Gln Ala Ile Arg Ile Arg Thr Gly Glu Thr Gly Glu Asp Ala Val  
100 105 110

## (2) INFORMATION FOR SEQ ID NO:9:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 112 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met Lys Lys Ile Glu Ala Ile Ile Arg Pro Phe Lys Leu Asp Glu Val  
1 5 10 15

Lys Ile Ala Leu Val Asn Ala Gly Ile Val Gly Met Thr Val Ser Glu  
20 25 30

Val Arg Gly Phe Gly Arg Gln Lys Gly Gln Thr Glu Arg Tyr Arg Gly  
35 40 45

Ser Glu Tyr Thr Val Glu Phe Leu Gln Lys Leu Lys Leu Glu Ile Val  
50 55 60

Val Glu Asp Ala Gln Val Asp Thr Val Ile Asp Lys Ile Val Ala Ala  
65 70 75 80

Ala Arg Thr Gly Glu Ile Gly Asp Gly Lys Ile Phe Val Ser Pro Val  
85 90 95

Asp Gln Thr Ile Arg Ile Arg Thr Gly Glu Lys Asn Ala Asp Ala Ile  
100 105 110

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 105 amino acids

-continued

(B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Lys Met Ile Lys Ala Ile Val Arg Pro Asp Lys Val Asp Asp Ile  
 1 5 10 15  
 Val Asp Ser Leu Glu Asn Ala Gly Tyr Pro Ala Phe Thr Lys Ile Asn  
 20 25 30  
 Ser Val Gly Arg Gly Lys Gln Gly Gly Leu Lys Val Gly Glu Ile Phe  
 35 40 45  
 Tyr Asp Glu Leu Pro Lys Thr Ile Leu Leu Ile Ala Val Asn Asp Asp  
 50 55 60  
 Glu Val Asp Glu Val Val Gly Leu Ile Lys Ser Ser Ala Ser Thr Gly  
 65 70 75 80  
 Asn Phe Gly Asp Gly Lys Ile Phe Ile Gln Pro Ile Thr Glu Ala Tyr  
 85 90 95  
 Thr Ile Arg Thr Gly Glu Thr Gly Ile  
 100 105

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 111 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met Lys Glu Val Ile Ala Ile Ile Arg Pro Asn Thr Val Ser Lys Thr  
 1 5 10 15  
 Val Lys Ala Leu Asp Val Val Gly Phe Pro Ala Val Thr Met Ala Glu  
 20 25 30  
 Cys Phe Gly Arg Gly Lys Gln Lys Gly Tyr Glu Glu Gly Glu Lys Glu  
 35 40 45  
 Gly Arg Phe Ile Lys Tyr Ile Pro Lys Arg Leu Ile Ser Ile Val Val  
 50 55 60  
 Asp Asp Ala Asp Val Pro Leu Val Val Gly Ile Ile Ser Lys Val Asn  
 65 70 75 80  
 Arg Thr Gly Ser Phe Gly Asp Gly Arg Ile Phe Val Leu Pro Val Glu  
 85 90 95  
 Glu Ala Ile Arg Val Arg Thr Gly Glu Thr Gly Glu Ile Ala Ile  
 100 105 110

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 15 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Phe Ser Ala Asn Leu Pro Glu Ile Val Asp Ile Gln Lys Ile Ile  
 1 5 10 15

-continued

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 817 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

```

CTGAAAGTTG TGTAAAAA AAAACTAGAA TCATGGCGGC GTCAATGACG AAACCCATCT 60
CAATAACTTC TCTCGGTTTC TATTCTGATC GAAAGACAT TGCTTTCTCT GATTGCATTT 120
CGATTTGTTC TGGATTGAGA CATTCCCAC CATCTTGCCCT CGATTTGGTC ACAAAGTCAC 180
CGAGTAATAA CAGTCGTGTT TTACCTGTGC TTAGTGCCCA AATATCTTCT GATTATATTC 240
CAGACTCGAA ATTTTACAAG GTGGAAGCAA TTGTCAGACC ATGGAGAATC CAGCAAGTTT 300
CATCGGCTTT ACTGAAAATC GGGATTGCGAG GTGTTACTGT TTCTGATGTG AGAGGGTTTG 360
GTGCACAAGG AGGTTCTACC GAGAGACACG GTGGCTCTGA GTTCTCGGAA GACAAAATTTG 420
TTGCTAAAGT TAAGATGGAA ATCGTTGTTA AGAAAGACCA AGTGGAACTCT GTAATCAACA 480
CAATAAATGA AGGAGCAAGG ACAGGAGAGA TTGGTGATGG CAAGATTTTT GTTTTGCCTG 540
TGTCAGATGT CATAAGAGTT AGGACAGGTG AGCGTGGGGA GAAAGCAGAG AAGATGACTG 600
GTGATATGCT TTCACCTCT TAGGAACAAA CAGAGCTCAA GAATGGTTTT TTTTTTTTTC 660
ATTTGCTCT CTAGATCTCG CGAATAATAA TGAATGGAGT CTGTGTTTG TTTTATGTTG 720
AATCGATCAA GATGTGTTTT TAACTGTACA TGAATTATGC AGAAACATCT GTCCTGGTTC 780
TCAGACATCG AACTCTGTT CTAATAAAAA AAAAAA 817

```

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 897 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: unknown  
 (D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

```

GCGGTGTCGG CCGCTCTAGA ACTAGTGGAT CCCC CGGCT GCAGGAATTC GGCACGAGGC 60
TACTGCGAAA CTGGGCTTGC TCACTCCTCT TCATTTCTAAT AACATCAAGA AAGAATTTCC 120
TGTTTTTGAT TTCAGTTTGT TTTGTCCAGA GCTTAGACAT TCTCGTTTTT CTCACTTTAA 180
CACCGCGGTC AAGCGCGTAA GATATGCCCC CGTCGTTTCT GTGATTAATG CCCAAAGCTC 240
GCCTGACTAC ATTCTGATG CTAAATTTCTA CAAAGTGAA GCAATTCTCA GGCCTGGCG 300
AGTCTCGCAA GTTTCTCGG CTTTGCTAAA AATTGGTATT CGAGGTGTTA CTGTTTCTGA 360
TGTTGAGGT TTTGGTGCT AAGGTGGTTC AACTGAGAGG CAGGGCGGCT CAGAAATTTT 420
TGAAGACAAG TTTGTTGCTA AAGTTAAGAT GGAGATCGTG GTTAGCAAAG ACCAGGTTGA 480
GGATGTTATA GAAAAATCA TTGAGGAGGC AAGAACTGGA GAGATTGGAG ACGGCAAGAT 540
TTCTTGCTG CCTGTTTCAG ATGTAATAAG AGTCCGCACT GGTGAGCGGG GTGATAAGGC 600
TGAGAGGATG ACAGGAGGGC GATCTGACAT GAGTACTTCT GCTTGACTGC TGTGACCAGC 660
AATATAGCAT TCAGACTAA CTGTCCTTTG AGAAAGCCCC GCCCTTATTA GCCATTATCC 720
AGTATAGCTT GATAATTGA ATTTTTTGTT TTCCTAATA AAGAAACAAA GATCTTTTCA 780

```



-continued

---

TTATCCTGTT GATGATAATT GAAAACGGAA GGATCGCGAA TTTGTTCAAG TGCTTGCAAG 840  
ATAAATAACA AGAAGAGGAG TAATGTTAAC AAAAAAAAAA AAAAAAAAAA ACTCGAG 897

## (2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 588 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATGGCGCGCT CAATGACGAA ACCCATCTCA ATAACCTCTC TCGGTTTCTA TTCTGATCGA 60  
 AAGAACATGG CTTTCTCTGA TTGCATTTTCG ATTTGTTCTG GATTCAGACA TTCCCGACCA 120  
 TCTTGCCCTCG ATTTGGTCAC AAAGTCACCG AGTAATAACA GTCGTGTTTT ACCTGTCGTT 180  
 AGTGCCCAA TATCTTCTGA TTATATTCCA GACTCGAAAT TTTACAAGGT GGAAGCAATT 240  
 GTCAGACCAT GGAGAATCCA GCAAGTTTCA TCGGCTTTAC TGAAAATCGG GATTCGAGGT 300  
 GTTACTGTTT CTGATGTGAG AGGGTTTGGT GCACAAGGAG GTTCTACCGA GAGACACGGT 360  
 GGCTCTGAGT TCTCGGAAGA CAAATTTGTT GCTAAAGTTA AGATGGAAAT CGTTGTTAAG 420  
 AAAGACCAAG TGGAAATCTGT AATCAACACA ATAATTGAAG GAGCAAGGAC AGGAGAGATT 480  
 GGTGATGGCA AGATTTTTGT TTTGCCTGTG TCAGATGTCA TAAGAGTTAG GACAGGTGAG 540  
 CGTGGGGAGA AAGCAGAGAA GATGACTGGT GATATGCTTT CACCGTCT 588

## (2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 594 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CGGCACGAGG CTACTGCGAA ACTGGGCTTG CTCACTCCTC TTCATTCTAA TAACATCAAG 60  
 AAAGAATTCC CTGTTTTTGA TTTCAGTTTG TTTTGTCCAG AGCTTAGACA TTCTCGGTTT 120  
 TCTCACTTTA ACACCGCGGT CAAGCGCGTA AGATATGCCC CCGTCGTTCC TGTGATTAAT 180  
 GCCCAAAGCT CGCCTGACTA CATTCTGAT GCTAAATCTT ACAAAGTGGA AGCAATTCTC 240  
 AGGCCCTGGC GAGTCTCGCA AGTTTCTCG GCTTTGCTAA AAATTGGTAT TCGAGGTGTT 300  
 ACTGTTTCTG ATGTTTCGAG TTTTGGTGCT CAAGTGTTT CAACTGAGAG GCAGGGCGGC 360  
 TCAGAATTTT CTGAAGACAA GTTTGTTGCT AAAGTTAAGA TGGAGATCGT GGTAGCAAA 420  
 GACCAGGTTG AGGATGTTAT AGAAAAAATC ATTGAGGAGG CAAGAAGTGG AGAGATTGGA 480  
 GACGGCAAGA TTTTCTTGTCT GCCTGTTTCA GATGTAATAA GAGTCCGCAC TGGTGAGCGG 540  
 GGTGATAAGG CTGAGAGGAT GACAGGAGGG CGATCTGACA TGAGTACTTC TGCT 594

## (2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 20 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

-continued

---

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCAAGATGGT CGGGAATGTC 20

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CGACAGGTAA AACACGACTG 20

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 20 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GGTCTGACAA TTGCTTCCAC 20

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GAAACCAAAC ACAGACTCC 19

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CCGAGTAATA ACAGTCGTC 19

---

What is claimed is:

1. A substantially pure nucleic acid molecule comprising a nucleotide sequence that encodes a P-II gene product comprising:

(a) the amino acid sequence of SEQ ID NO:1; or

(b) the amino acid sequence of SEQ ID NO:2.

2. A substantially pure nucleic acid molecule encoding a P-II gene product, wherein said substantially pure nucleic acid molecule comprises:

(a) the nucleotide sequence of SEQ ID NO: 13;

(b) the nucleotide sequence of SEQ ID NO: 14;

(c) the nucleotide sequence of SEQ ID NO: 15; or

(d) the nucleotide sequence of SEQ ID NO: 16.

3. A substantially pure nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule of claim 1 under highly stringent conditions, wherein said substantially pure nucleic acid molecule encodes a functionally equivalent P-II gene product.

41

4. A substantially pure nucleic acid molecule which hybridizes to the complement of the nucleic acid molecule of claim 2 under highly stringent conditions, wherein said substantially pure nucleic acid molecule encodes a functionally equivalent P-PII gene product.

5. A vector comprising the substantially pure nucleic acid molecule as in any one of claims 1-4.

6. An expression vector comprising the substantially pure nucleic acid molecule as in any one of claims 1-4 operatively associated with a regulatory nucleotide sequence controlling the expression of the substantially pure nucleic acid molecule in a host cell.

7. A host cell genetically engineered to contain the substantially pure nucleic acid molecule as in any one of claims 1-4.

8. A host cell genetically engineered to express the substantially pure nucleic acid molecule as in any one of

42

claims 1-4 operatively associated with a regulatory nucleotide sequence controlling the expression of the substantially pure nucleic acid molecule in said host cell.

9. A transgenic plant containing a transgene comprising the substantially pure nucleic acid molecule as in any one of claims 1-4.

10. A transgenic plant containing a transgene comprising the substantially pure nucleic acid molecule as in any one of claims 1-4 operatively associated with a regulatory nucleotide sequence controlling the expression of the substantially pure nucleic acid molecule in a transgenic plant cell.

11. The transgenic plant of claim 9, wherein said transgene encodes an antisense nucleic acid molecule that suppresses expression of endogenous P-PII gene product.

\* \* \* \* \*